

## New England Biolabs Certificate of Analysis

**Product Name:** BgIII  
**Catalog Number:** R0144L  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in NEBuffer r3.1 in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10183327  
**Expiration Date:** 04/2025  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM TES, 500 mM NaCl, 200 µg/ml rAlbumin, 50% Glycerol, (pH 8.0 @ 25°C)  
**Specification Version:** PS-R0144S/L/E v3.0


| BgIII Component List |                              |            |                      |
|----------------------|------------------------------|------------|----------------------|
| NEB Part Number      | Component Description        | Lot Number | Individual QC Result |
| R0144LVIAL           | BgIII                        | 10183324   | Pass                 |
| B7024AVIAL           | Gel Loading Dye, Purple (6X) | 10182168   | Pass                 |
| B6003SVIAL           | NEBuffer™ r3.1               | 10182163   | Pass                 |

| Assay Name/Specification  | Lot # 10183327 |
|---|----------------|
| <b>Blue-White Screening (Terminal Integrity)</b><br>A sample of LITMUS28i vector linearized with a 10-fold excess of BgIII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.  | Pass           |
| <b>Endonuclease Activity (Nicking)</b><br>A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of BgIII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.            | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of BgIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass           |
| <b>Functional Testing (15 minute Digest)</b><br>A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and 1 µl of BgIII  | Pass           |

| Assay Name/Specification   | Lot # 10183327 |
|--|----------------|
| incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.   |                |
| <p><b>Ligation and Recutting (Terminal Integrity)</b><br/>After a 20-fold over-digestion of Lambda DNA with BglII, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with BglII.</p>  | <b>Pass</b>    |
| <p><b>Non-Specific DNase Activity (16 Hour)</b><br/>A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and a minimum of 100 units of BglII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>   | <b>Pass</b>    |
| <p><b>Protein Purity Assay (SDS-PAGE)</b><br/>BglII is &gt;95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>   | <b>Pass</b>    |
| <p><b>qPCR DNA Contamination (E. coli Genomic)</b><br/>A minimum of 10 units of BglII is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p> | <b>Pass</b>    |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



YunJie Sun  
Production Scientist  
30 Mar 2023



Michael Tonello  
Packaging Quality Control Inspector  
21 Apr 2023