

## New England Biolabs Certificate of Analysis

**Product Name:** BstBI  
**Catalog Number:** R0519S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 65°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10171829  
**Expiration Date:** 12/2024  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0519S/L v2.0

BstBI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0519SVIAL	BstBI	10171828	Pass
B6004SVIAL	rCutSmart™ Buffer	10168651	Pass

Assay Name/Specification	Lot # 10171829
<p><b>Non-Specific DNase Activity (16 Hour)</b>            A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 20 units of BstBI incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p><b>Ligation and Recutting (Terminal Integrity)</b>            After a 20-fold over-digestion of Lambda DNA with BstBI, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with BstBI.</p>	Pass
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 100 units of BstBI incubated for 4 hours at 65°C releases &lt;0.1% of the total radioactivity.</p>	Pass
<p><b>Functional Testing (15 minute Digest)</b>            A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of BstBI incubated for 15 minutes at 65°C results in complete digestion as determined</p>	Pass

Assay Name/Specification	Lot # 10171829
<p>by agarose gel electrophoresis.</p> <p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled pUC19 DNA and a minimum of 20 units of BstBI incubated for 4 hours at 65°C results in &lt;20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<p><b>Pass</b></p>


This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.




---

YunJie Sun  
Production Scientist  
08 Dec 2022




---

Michael Tonello  
Packaging Quality Control Inspector  
09 Dec 2022