

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: DpnII
Catalog Number: R0543S
Concentration: 10,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 μg

of Lambda DNA (dam-) in 1 hour at 37°C in a total reaction volume of

50 μl.

Lot Number: 10055674
Expiration Date: 08/2021
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R0543S/L v1.0

DpnII Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0543SVIAL	DpnII	10051234	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10050274	Pass	
B0543SVIAL	NEBuffer™ DpnII	10050225	Pass	

Assay Name/Specification	Lot # 10055674
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer DpnII containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 Units of DpnII incubated for 4 hours at 37°C results in <10%	Pass
conversion to the nicked form as determined by agarose gel electrophoresis.	_
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer DpnII containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of DpnII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda dam- DNA with DpnII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with DpnII.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer DpnII containing 1 µg of Lambda dam- DNA and a minimum	Pass



R0543S / Lot: 10055674

Page 1 of 2

Assay Name/Specification	Lot # 10055674
of 100 units of DpnII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) DpnII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Anthony Francis
Production Scientist

01 Aug 2019

Jay Minichiello

Packaging Quality Control Inspector

25 Sep 2019



R0543S / Lot: 10055674

Page 2 of 2