

## New England Biolabs Certificate of Analysis

*Product Name:* Cac8I  
*Catalog #:* R0579S/L  
*Concentration:* 5,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
*Lot #:* 0271603  
*Assay Date:* 03/2016  
*Expiration Date:* 3/2018  
*Storage Temp:* -20°C  
*Storage Conditions:* 150 mM KCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 0.10 % TritonX-100  
*Specification Version:* PS-R0579S/L v2.0  
*Effective Date:* 04 Aug 2015

Assay Name/Specification (minimum release criteria)	Lot #0271603
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 15 units of Cac8I incubated for 4 hours at 37°C releases <0.2% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 5-fold over-digestion of Lambda DNA with Cac8I, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 25°C. Of these ligated fragments, >95% can be recut with Cac8I.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 25 units of Cac8I incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>



Authorized by  
Derek Robinson  
04 Aug 2015



Inspected by  
Casey Madinger  
15 Mar 2016

