

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	PaqCl®
Catalog Number:	R0745S
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in rCutSmart Buffer incubated for 1 hour at 37°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10238214
Expiration Date:	10/2025
Storage Temperature:	-20°C
Storage Conditions:	50 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 μg/ml rAlbumin, (pH 7.4 @ 25°C)
Specification Version:	PS-R0745S/L v2.0

PaqCI® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
S0532SVIAL	PaqCI® Activator	10238216	Pass	
R0745SVIAL	PaqCl®	10230757	Pass	
B6004SVIAL	rCutSmart™ Buffer	10233338	Pass	

Assay Name/Specification	Lot # 10238214
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 units of PaqCI® incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of PaqCI® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with PaqCI® , >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PaqCI® .	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	Pass





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350

www.neb.com info@neb.com

Assay Name/Specification	Lot # 10238214
30 units of PaqCI® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) PaqCl® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 08 Apr 2024

Monkieuncy

Talia Monkiewicz Packaging Quality Control Inspector 08 Apr 2024

