

New England Biolabs Certificate of Analysis

Product Name: Nsil-HF[®]
Catalog Number: R3127S
Concentration: 20,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10206728
Expiration Date: 01/2025
Storage Temperature: -20°C
Storage Conditions: 300 mM NaCl , 10 mM Tris-HCl (pH 7.4), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 500 µg/ml BSA
Specification Version: PS-R3127S/L v1.0


| Nsil-HF [®] Component List | | | |
|-------------------------------------|-------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R3127SVIAL | Nsil-HF [®] | 10176640 | Pass |
| B7024AVIAL | Gel Loading Dye, Purple (6X) | 10198639 | Pass |
| B6004SVIAL | rCutSmart [™] Buffer | 10198644 | Pass |

| Assay Name/Specification | Lot # 10206728 |
|---|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of supercoiled φX174 DNA and a minimum of 100 units of Nsil-HF incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of Nsil-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Functional Test (15 minute Digest) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of Lambda DNA and 1 µl of Nsil-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with Nsil-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, | Pass |

| Assay Name/Specification | Lot # 10206728 |
|--|----------------|
| >95% can be recut with Nsil-HF. | |
| <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of Nsil-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | Pass |
| <p>Protein Purity Assay (SDS-PAGE) Nsil-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p> | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



YunJie Sun
Production Scientist
17 Jan 2023



Michael Tonello
Packaging Quality Control Inspector
05 Sep 2023