

DISCOVERY 240 County Road
GENUINE Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Bsal-HF®v2
Catalog Number: R3733L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number: 10040994
Expiration Date: 12/2020
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA,

50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-R3733S/L v1.0

Bsal-HF®v2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3733LVIAL	Bsal-HF®v2	10032391	Pass	
B7204SVIAL	CutSmart® Buffer	10036666	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10038708	Pass	

Assay Name/Specification	Lot # 10040994
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and	Pass
a minimum of 20 units of Bsal-HF®v2 incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Bsal-HF®v2 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Protein Purity Assay (SDS-PAGE) Bsal-HF®v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of pXba DNA and a minimum of 60 units of Bsal-HF®v2 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



R3733L / Lot: 10040994

Page 1 of 2



Assay Name/Specification	Lot # 10040994
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with Bsal-HF®v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Bsal-HF®v2.	Pass

This product has been tested and shown to be in compliance with all specifications.

Anthony Francis
Production Scientist

27 Dec 2018

Michael Tonello

Packaging Quality Control Inspector

08 Apr 2019

