

## New England Biolabs Product Specification

*Product Name:* SwaI  
*Catalog #:* R0604S/L  
*Concentration:* 10,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 25°C in a total reaction volume of 50 µl.  
*Shelf Life:* 24 months  
*Storage Temp:* -20°C  
*Storage Conditions:* 400 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
*Specification Version:* PS-R0604S/L v2.0  
*Effective Date:* 28 Mar 2016

### Assay Name/Specification (minimum release criteria)

**Exonuclease Activity (Radioactivity Release)** - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 100 units of SwaI incubated for 4 hours at 25°C releases <0.1% of the total radioactivity.

**Ligation and Recutting (Terminal Integrity)** - After a 20-fold over-digestion of pXba-NdeI DNA with SwaI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with SwaI.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of pXba DNA and a minimum of 100 units of SwaI incubated for 16 hours at 25°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Date 28 Mar 2016

Derek Robinson  
Director of Quality Control

